arXiv:1601.07096v1 [math.CT] 26 Jan 2016

Group-groupoid actions and liftings of crossed modules

Osman Mucuk*[a] and Tunçar Şahan†[b]

[a]Department of Mathematics, Erciyes University, Kayseri, TURKEY
[b]Department of Mathematics, Aksaray University, Aksaray, TURKEY

Abstract

The aim of this paper is to define the notion of lifting of a crossed module via a group morphism and give some properties of this type of the lifting. Further we obtain a criterion for a crossed module to have a lifting of crossed module. We also prove that the liftings of a certain crossed module constitute a category; and that this category is equivalent to the category of covers of that crossed module and hence to the category of group-groupoid actions of the corresponding groupoid to that crossed module.

Key Words: Crossed module, covering, lifting, action groupoid
Classification: 20L05, 57M10, 22AXX, 22A22, 18D35

1 Introduction

The theory of covering groupoids has an important role in the applications of groupoids (cf. [2] and [12]). In this theory it is well known that for a groupoid G, the category $\mathsf{GpdAct}(\mathsf{G})$ of groupoid actions of G on sets, these are also called operations or G-sets, are equivalent to the category $\mathsf{GpdCov}/\mathsf{G}$ of covering groupoids of G (see [5, Theorem 2] for the topological version of this equivalence).

On the one hand in [6, Proposition 3.1] it was proved that if G is a group-groupoid, which is an internal groupoid in the category of groups and widely used in literature under the names *2-group* (see for example [3]), *$\mathcal{G}$-groupoid* or *group object* [7] in the category of

*E-mail : mucuk@erciyes.edu.tr
†E-mail : tuncarsahan@aksaray.edu.tr

groupoids, then the category $\mathsf{GpGpdCov/G}$ of group-groupoid coverings of G is equivalent to the category $\mathsf{GpGpdAct(G)}$ of group-groupoid actions of G on groups. In [1, Theorem 4.2] this result has been recently generalized to the case where G is an internal groupoid for an algebraic category C which includes groups, rings without identity, R-modules, Lie algebras, Jordan algebras, and many others; acting on a group with operations in the sense of Orzech [18].

On the other hand in [7, Theorem 1] it was proved that the categories of crossed modules and group-groupoids, under the name of $\mathcal{G}$-groupoids, are equivalent (see also [13] for an alternative equivalence in terms of an algebraic object called *cat^n-groups*). By applying this equivalence of the categories, normal and quotient objects in the category of group-groupoids have been recently obtained in [16]. In [19, Section 3] it was proved that a similar result to the one in [7, Theorem 1] can be generalized for a certain algebraic category introduced by Orzech [18], and adapted and called category of groups with operations. The study of internal category theory was continued in the works of Datuashvili [10] and [11]. Moreover, she developed cohomology theory of internal categories, equivalently, crossed modules, in categories of groups with operations [8] and [9]. The equivalences of the categories in [7, Theorem 1] and [19, Section 3] enable us to generalize some results on group-groupoids to the more general internal groupoids for a certain algebraic category C (see for example [1], [14], [15] and [17]).

In this paper we use the equivalence of group-groupoids and crossed modules proved in [7, Theorem 1] to determine a notion in crossed modules, called lifting of a crossed module, corresponding to a group-groupoid action, investigate some properties of lifting of crossed modules and give a criteria for a crossed module to have a lifting. Finally we prove that for a group-groupoid G, the group-groupoid actions of G on groups and liftings of the crossed module corresponding to G are categorically equivalent.

2 Preliminaries

Let G be a groupoid. We write $\mathrm{Ob}(G)$ for the set of objects of G and write G for the set of morphisms. We also identify $\mathrm{Ob}(G)$ with the set of identities of G and so an element of $\mathrm{Ob}(G)$ may be written as x or 1_x as convenient. We write $d_0, d_1 \colon G \to \mathrm{Ob}(G)$ for the source and target maps, and, as usual, write $G(x, y)$ for $d_0^{-1}(x) \cap d_1^{-1}(y)$, for $x, y \in \mathrm{Ob}(G)$. The composition $h \circ g$ of two elements of G is defined if and only if $d_0(h) = d_1(g)$, and so the map $(h, g) \mapsto h \circ g$ is defined on the pullback $G_{d_0} \times_{d_1} G$ of d_0 and d_1. The *inverse* of $g \in G(x, y)$ is denoted by $g^{-1} \in G(y, x)$.

If $x \in \mathrm{Ob}(G)$, we write $\mathsf{St}_G x$ for $d_0^{-1}(x)$ and call the *star* of G at x. Similarly we write

$\mathsf{Cost}_G x$ for $d_1^{-1}(x)$ and call *costar* of G at x. The set of all morphisms from x to x is a group, called *object group* at x, and denoted by $G(x)$.

A groupoid G is *transitive (resp. simply transitive, 1-transitive and totally intransitive)* if $G(x,y) \neq \emptyset$ (resp. $G(x,y)$ has no more than one element, $G(x,y)$ has exactly one element and $G(x,y) = \emptyset$) for all $x, y \in \mathrm{Ob}(G)$ such that $x \neq y$.

A totally intransitive groupoid is determined entirely by the family $\{G(x) \mid x \in \mathrm{Ob}(G)\}$ of groups. This totally intransitive groupoid is sometimes called *totally disconnected* or *bundle of groups* [2, pp.218].

Let $p\colon \widetilde{G} \to G$ be a morphism of groupoids. Then p is called a *covering morphism* and $\widetilde{G}$ a *covering groupoid* of G if for each $\widetilde{x} \in \mathrm{Ob}(\widetilde{G})$ the restriction $\mathsf{St}_{\widetilde{G}}\widetilde{x} \to \mathsf{St}_G p(\widetilde{x})$ is bijective.

Assume that $p\colon \widetilde{G} \to G$ is a covering morphism. Then we have a lifting function $S_p\colon G_{d_0} \times_{\mathrm{Ob}(p)} \mathrm{Ob}(\widetilde{G}) \to \widetilde{G}$ assigning to the pair (a,x) in the pullback $G_{d_0} \times_{\mathrm{Ob}(p)} \mathrm{Ob}(\widetilde{G})$ the unique element b of $\mathsf{St}_{\widetilde{G}} x$ such that $p(b) = a$. Clearly S_p is inverse to $(p, d_0)\colon \widetilde{G} \to G_s \times_{\mathrm{Ob}(p)} \mathrm{Ob}(\widetilde{G})$. So it is stated that $p\colon \widetilde{G} \to G$ is a covering morphism if and only if (p,s) is a bijection.

A covering morphism $p\colon \widetilde{G} \to G$ is called *transitive* if both $\widetilde{G}$ and G are transitive. A transitive covering morphism $p\colon \widetilde{G} \to G$ is called *universal* if $\widetilde{G}$ covers every cover of G, i.e., for every covering morphism $q\colon \widetilde{H} \to G$ there is a unique morphism of groupoids $\widetilde{p}\colon \widetilde{G} \to \widetilde{H}$ such that $q\widetilde{p} = p$ (and hence $\widetilde{p}$ is also a covering morphism), this is equivalent to that for $\widetilde{x}, \widetilde{y} \in \mathrm{Ob}(\widetilde{G})$ the set $\widetilde{G}(\widetilde{x}, \widetilde{y})$ has not more than one element.

Recall that an action of a groupoid G on a set S via a function $\omega\colon S \to \mathrm{Ob}(G)$ is a function $G_{d_0} \times_\omega S \to S, (g,s) \mapsto g \bullet s$ satisfying the usual rules for an action: $\omega(g \bullet s) = d_1(g)$, $1_{\omega(s)} \bullet s = s$ and $(h \circ g) \bullet s = h \bullet (g \bullet s)$ whenever $h \circ g$ and $g \bullet s$ are defined. A morphism $f\colon (S,\omega) \to (S',\omega')$ of such actions is a function $f\colon S \to S'$ such that $w' f = w$ and $f(g \bullet s) = g \bullet f(s)$ whenever $g \bullet s$ is defined. This gives a category $\mathsf{GpdAct}(\mathsf{G})$ of actions of G on sets. For such an action the action groupoid $G \ltimes S$ is defined to have object set S, morphisms the pairs (g,s) such that $d_0(g) = \omega(s)$, source and target maps $d_0(g,s) = s$, $d_1(g,s) = g \bullet s$, and the composition

$$(g', s') \circ (g, s) = (g \circ g', s)$$

whenever $s' = g \bullet s$. The projection $q\colon G \ltimes S \to G, (g,s) \mapsto s$ is a covering morphism of groupoids and the functor assigning this covering morphism to an action gives an equivalence of the categories $\mathsf{GpdAct}(\mathsf{G})$ and $\mathsf{GpdCov}/\mathsf{G}$.

3 Group-groupoids and crossed modules

A *group-groupoid* is a groupoid G with morphisms of groupoids $G \times G \to G$, $(g, h) \mapsto g+h$ and $G \to G, g \to -g$ yielding a group structure internal to the category of groupoids. Since the addition map is a morphism of groupoids, we have an interchange rule that $(b \circ a) + (d \circ c) = (b + d) \circ (a + c)$ for all $a, b, c, d \in G$ such that $b \circ a$ and $d \circ c$ are defined. If the identity of $\mathrm{Ob}(G)$ is e, then the identity of the group structure on the morphisms set is 1_e.

Let G be a group-groupoid. An action of the group-groupoid G on a group X via ω consists of a morphism $\omega \colon X \to \mathrm{Ob}(G)$ from the group X to the underlying group of $\mathrm{Ob}(G)$ and an action of the groupoid G on the underlying set X via ω such that the following interchange law holds:

$$(g \bullet x) + (g' \bullet x') = (g + g') \bullet (x + x') \tag{1}$$

whenever both sides are defined. A morphism $f \colon (X, \omega) \to (X', \omega')$ of such actions is a morphism $f \colon X \to X'$ of groups and of the underlying operations of G. This gives a category $\mathsf{GpGpdAct(G)}$ of actions of G on groups. For an action of G on the group X via ω, the action groupoid $G \ltimes X$ has a group structure defined by

$$(g, x) + (g', x') = (g + g', x + x')$$

and with this operation $G \ltimes X$ becomes a group-groupoid and the projection $p \colon G \ltimes X \to G$ is an object of the category $\mathsf{GpGpdCov/G}$. By means of this construction the following equivalence of the categories was given in [6, Proposition 3.1].

Proposition 3.1. *The categories* $\mathsf{GpGpdCov/G}$ *and* $\mathsf{GpGpdAct(G)}$ *are equivalent.*

We recall that as defined by Whitehead in [20, 22] a crossed module of groups consists of two groups A and B, an action of B on A denoted by $b \cdot a$ for $a \in A$ and $b \in B$; and a morphism $\alpha \colon A \to B$ of groups satisfying the following conditions for all $a, a_1 \in A$ and $b \in B$

CM1 $\alpha(b \cdot a) = b + \alpha(a) - b$,

CM2 $\alpha(a) \cdot a_1 = a + a_1 - a$.

We will denote such a crossed module by (A, B, α).

Here are some examples of well known crossed modules:

(i) The inclusion map $N \hookrightarrow G$ of a normal subgroup is a crossed module with the conjugation action of G on N.

(ii) If M is a G-module, then the zero map $0\colon M \to G$ has the structure of crossed module.

(iii) The inner automorphism map $G \to Aut(G)$ is a crossed module.

(iv) If X is a topological group, then the fundamental group πX is a group-groupoid, the star $\mathsf{St}_{\pi X}0$ at the identity $0 \in X$ becomes a group and the final point map $d_1\colon \mathsf{St}_{\pi X}0 \to X$ becomes a crossed module.

(v) As a motivating geometric example of crossed module due to Whitehead [20, 22] if X is topological space and $A \subseteq X$ with $x \in A$, then there is a natural action of $\pi_1(A, x)$ on second relative homotopy group $\pi_2(X, A, x)$ and with this action the boundary map

$$\partial\colon \pi_2(X, A, x) \to \pi_1(A, x)$$

becomes a crossed module. This crossed module is called *fundamental crossed module* and denoted by $\Pi(X, A, x)$ (see for example [4] for more details).

The following are some standard properties of crossed modules.

Proposition 3.2. *Let (A, B, α) be a crossed module. Then*

(i) $\alpha(A)$ is a normal subgroup of B.

(ii) $\operatorname{Ker}\alpha$ is central in A, i.e. $\operatorname{Ker}\alpha$ is a subset of $Z(A)$, the center of A.

(iii) $\alpha(A)$ acts trivially on $Z(A)$.

(iv) $Z(A)$ and $\operatorname{Ker}\alpha$ inherit an action of $\operatorname{Coker}\alpha$ to become $(\operatorname{Coker}\alpha)$-modules.

Let (A, B, α) and (A', B', α') be two crossed modules. A morphism (f_1, f_2) from (A, B, α) to (A', B', α') is a pair of morphisms of groups $f_1\colon A \to A'$ and $f_2\colon B \to B'$ such that $f_2\alpha = \alpha' f_1$ and $f_1(b \cdot a) = f_2(b) \cdot f_1(a)$ for $a \in A$ and $b \in B$.

Crossed modules with morphisms between them form a category denoted by XMod.

The following result was proved in [7, Theorem 1] and since we need some details of the proof, we give a sketch proof of the equivalence.

Theorem 3.1. *The category* XMod *of crossed modules and the category* GrpGpd *of group-groupoids are equivalent.*

Proof. Let GrpGpd be the category of group-groupoids and XMod the category of crossed modules.

A functor $\delta\colon \mathsf{GrpGpd} \to \mathsf{XMod}$ is defined as follows. For a group-groupoid G let $\delta(G)$ be the crossed module (A, B, d_1) where $A = \operatorname{Ker} d_0$, $B = \operatorname{Ob}(G)$ and $d_1\colon A \to B$ is the restriction of the target point map . Then A, B inherit group structures from that of G, and the target point map $d_1\colon A \to B$ is a morphism of groups. Further we have an action $B \times A \to A$, $(b, a) \mapsto b \cdot a$ of B on the group A given by

$$b \cdot a = 1_b + a - 1_b \tag{2}$$

for $a \in A$, $b \in B$ and we clearly have

$$d_1(b \cdot a) = b + d_1(a) - b \tag{3}$$

and

$$d_1(a) \cdot a_1 = a + a_1 - a \tag{4}$$

for $a, a_1 \in A$, $b \in B$. Thus (A, B, d_1) is a crossed module.

Conversely define a functor $\eta\colon \mathsf{XMod} \to \mathsf{GrpGpd}$ in the following way. For a crossed module (A, B, α) let $\eta(A, B, \alpha)$ be the group-groupoid whose object set (group) is the group B and whose group of the morphisms is the semi-direct group $A \rtimes B$ with the usual group structure

$$(a_1, b_1) + (a, b) = (a_1 + b_1 \cdot a, b_1 + b) \tag{5}$$

The source and target point maps are defined to be $d_0(a, b) = b$ and $d_1(a, b) = \alpha(a) + b$, while the groupoid composition is given by

$$(a_1, b_1) \circ (a, b) = (a_1 + a, b)$$

whenever $b_1 = \alpha(a) + b$. □

Proposition 3.3. *Let G be a group-groupoid and (A, B, α) the crossed module corresponding to G. If G is transitive (resp. simply transitive, 1-transitive and totally intransitive) then α is surjective (resp. injective, bijective; and zero morphism such that A is abelian).*

Proof. The proof can be followed by the details of the proof of Theorem 3.1. □

Hence we can state the following definition.

Definition 3.1. Let (A, B, α) be a crossed module. Then (A, B, α) is called *transitive (resp. simply transitive, 1-transitive and totally intransitive) if α is surjective (resp. injective, bijective; and zero morphism such that A is abelian).*

Example 3.1. If X is a topological group whose underlying topology is path-connected (resp. totally disconnected), then the crossed module $(\mathsf{St}_{\pi X}0, X, d_1)$ is transitive (resp. totally intransitive).

4 Liftings of crossed modules

In this section using Theorem 3.1, we determine the notion in crossed modules, corresponding to the action of a group-groupoid on a group and interpret the properties in crossed modules corresponding to group-groupoid actions .

Let G be a group-groupoid acting on a group X by an action $G_{d_0} \times_\omega X, (g, x) \mapsto g \bullet x$, via a group morphism $\omega \colon X \to \mathrm{Ob}(G)$ and let (A, B, α) be the crossed module corresponding to G. Then we have a morphism $\omega \colon X \to B$ of groups and an action of X on $A = \mathsf{St}_G 0$ defined by

$$X \times A \to A, x \cdot a = 1_{\omega(x)} + a - 1_{\omega(x)} \tag{6}$$

By the group-groupoid action of G on X we have a group morphism

$$\varphi \colon A \to X, a \mapsto \varphi(a) = a \bullet 0_X \tag{7}$$

such that $\omega\varphi = \alpha$, where 0_X is the identity element of the group X.

Then we prove the following theorems.

Theorem 4.1. *By the action of X on A defined above, (A, X, φ) becomes a crossed module.*

Proof. We prove that the conditions [CM1] and [CM2] are satisfied.

[CM1] For all $a \in A$ and $x \in X$ we have the following evaluations

$$\begin{aligned}
\varphi(x \cdot a) &= \varphi(1_{\omega(x)} + a - 1_{\omega(x)}) && \text{(by Eq. 6)} \\
&= (1_{\omega(x)} + a - 1_{\omega(x)}) \bullet 0_X && \text{(by Eq. 7)} \\
&= (1_{\omega(x)} + (a + 1_{\omega(-x)})) \bullet (x + (-x)) \\
&= (1_{\omega(x)} \bullet x) + ((a + 1_{\omega(-x)}) \bullet (-x)) && \text{(by Eq. 1)} \\
&= (0_A + 1_{\omega(x)}) \bullet (0_X + x) + (a + 1_{\omega(-x)}) \bullet (0_X + (-x))
\end{aligned}$$

and by the interchange law (1)

$$\begin{aligned}\varphi(x \cdot a) &= (0_A \bullet 0_X) + (1_{\omega(x)} \bullet x) + (a \bullet 0_X) + (1_{\omega(-x)} \bullet (-x)) \\ &= \varphi(0_A) + x + \varphi(a) - x && \text{(by Eq. 7)} \\ &= x + \varphi(a) - x\end{aligned}$$

[CM2] For all $a, a_1 \in A$ we have the following steps

$$\begin{aligned}\varphi(a) \cdot a_1 &= (a \bullet 0_X) \cdot a_1 && \text{(by Eq. 7)} \\ &= 1_{\omega(a \bullet 0_X)} + a_1 - 1_{\omega(a \bullet 0_X)} && \text{(by Eq. 6)} \\ &= 1_{d_1(a)} + a_1 - 1_{d_1(a)} && (\text{ by } \omega(a \bullet 0_X) = d_1(a)) \\ &= d_1(a) \cdot a_1 && \text{(by Eq. 2)} \\ &= a + a_1 - a && \text{(by Eq. 4)}\end{aligned}$$

□

Theorem 4.2. *Let (A, B, α) be a crossed module and $\omega\colon X \to B$ a group morphism. Then any group morphism $\varphi\colon A \to X$ such that $\omega\varphi = \alpha$ is a crossed module with the action defined via ω if and only if the map $\overline{\varphi}\colon A \rtimes X \to X$ defined by*

$$\overline{\varphi}(a, x) = \varphi(a) + x \tag{8}$$

is a group morphism.

Proof. Suppose that the map $\overline{\varphi}\colon A \rtimes X \to X$ defined by $\overline{\varphi}(a, x) = \varphi(a) + x$ is a group morphism. Then we have the following evaluations to prove that the axioms [CM1] and [CM2] of crossed module are satisfied for the group morphism $\varphi\colon A \to X$.

[CM1] For $x \in X$ and $a \in A$ we have that

$$\begin{aligned}\varphi(x \cdot a) = \overline{\varphi}(x \cdot a, 0) &= \overline{\varphi}((0, x) + (a, -x)) && \text{(by Eq. 5)} \\ &= \overline{\varphi}(0, x) + \overline{\varphi}(a, -x) \\ &= \varphi(0) + x + \varphi(a) - x && \text{(by Eq. 8)} \\ &= x + \varphi(a) - x\end{aligned}$$

[CM2] Since the action of X on A is defined by means of the action of B on A via ω, the morphism φ satisfies the condition **[CM2]** of crossed module.

Conversely assume that $\varphi\colon A \to X$ is a crossed module. Then by the following evaluation the map $\overline{\varphi}\colon A \rtimes X \to X$ becomes a group morphism.

$$\begin{aligned}
\overline{\varphi}((a,x)+(a_1,x_1)) &= \overline{\varphi}(a + x\cdot a_1, x + x_1) && \text{(by Eq. 5)}\\
&= \varphi(a + x\cdot a_1) + (x + x_1) && \text{(by Eq. 8)}\\
&= \varphi(a) + \varphi(x\cdot a_1) + (x + x_1)\\
&= \varphi(a) + x + \varphi(a_1) - x + x + x_1 && \text{(by \textbf{CM1})}\\
&= \varphi(a) + x + \varphi(a_1) + x_1\\
&= \overline{\varphi}(a,x) + \overline{\varphi}(a_1,x_1). && \text{(by Eq. 8)}
\end{aligned}$$

□

We now define the notion of lifting of a crossed module as follows:

Definition 4.1. Let (A, B, α) be a crossed module and $\omega\colon X \to B$ a morphism of groups. Then a crossed module (A, X, φ) in which the action of X on A is defined via ω, is a called a *lifting of* α *over* ω and denoted by (φ, X, ω) if the following diagram is commutative

$$\begin{array}{ccc}
 & & X \\
 & \overset{\varphi}{\nearrow} & \downarrow{\scriptstyle \omega} \\
A & \xrightarrow[\alpha]{} & B
\end{array}$$

It is obvious that every crossed module (A, B, α) lifts to itself over the identity morphism 1_B on B.

Remark 4.1. In the following diagram if (φ, X, ω) is a lifting of (A, B, α), then $\operatorname{Ker}\varphi \subseteq \operatorname{Ker}\alpha$ and $(1_A, \omega)$ is a morphism of crossed modules

$$\begin{array}{ccc}
A & \overset{\varphi}{\rightarrowtail} & X \\
{\scriptstyle 1_A}\| & & \downarrow{\scriptstyle \omega} \\
A & \xrightarrow[\alpha]{} & B
\end{array}$$

Therefore if (A, B, α) is a simply transitive crossed module then $\operatorname{Ker}\alpha$ is trivial and so also $\operatorname{Ker}\varphi$ is. Hence the crossed module (A, X, φ) is also simply transitive.

The following can be stated as examples of lifting crossed modules.

Example 4.1. A crossed module (A, B, α) is a lifting of the automorphism crossed module

$(A, \mathrm{Aut}(A), \iota)$ over the action of B on A, i.e. $\theta\colon B \to \mathrm{Aut}(A)$, $\theta(b)(a) = b \cdot a$



Example 4.2. Let N be a normal subgroup of a group G. Since (N, G, inc) is a simply transitive crossed module, any lifting crossed module (N, X, φ) is also simply transitive. Moreover $\varphi(N)$ is a normal subgroup of X. Since φ is injective N also can be consider as a normal subgroup of X.

Example 4.3. Let G be a group with trivial center. Then the automorphism crossed module $G \to \mathrm{Aut}\, G$ is simply transitive and hence every lifting of $G \to \mathrm{Aut}\, G$ is also simply transitive.

Example 4.4. Let $p\colon \widetilde{X} \to X$ be a covering morphism of topological groups. Hence $(\mathsf{St}_{\pi X}0, X, d_1)$ is a crossed module. If α is a path in X with initial point $0 \in X$, the identity, then by the path lifting property there exists a unique path $\widetilde{\alpha}$ in $\widetilde{X}$ such that $p\widetilde{\alpha} = \alpha$ and $\widetilde{\alpha}(0) = \tilde{0} \in \widetilde{X}$, the identity. Hence we can define a function $\widetilde{d_1}\colon \mathsf{St}_{\pi X}0 \to \widetilde{X}$ assigning $[\alpha] \in \mathsf{St}_{\pi X}0$ to the final point $\widetilde{\alpha}(1)$ of the lifting path of α at $\widetilde{0}$. It follows that $\widetilde{d_1}$ is well defined and $(\widetilde{d_1}, \widetilde{X}, p)$ is a lifting of $(\mathsf{St}_{\pi X}0, X, d_1)$.

We now state some results on lifting crossed modules.

Lemma 4.1. *Let (A, B, α) be a crossed module and φ a lifting of α over $\omega\colon X \to B$. If there are isomorphisms $f\colon B \to B'$ and $g\colon X' \to X$ for some groups B' and G' then φ' is a lifting of α' over $\omega'\colon X' \to B'$ where $\varphi' = g^{-1}\varphi$, $\alpha' = f\alpha$ and $\omega' = f\omega g$.*

Proof. It follows that $\alpha' = f\alpha$ and $\varphi' = g^{-1}\varphi$ are crossed modules since f and g are isomorphisms; and

$$\begin{aligned}\omega'\varphi' &= (f\omega g)(g^{-1}\varphi)\\ &= f\omega\varphi\end{aligned}$$

Since (φ, X, ω) is a lifting of (A, B, α), i.e., $\omega\varphi = \alpha$ we have that

$$\begin{aligned}\omega'\varphi' &= f\omega\varphi\\ &= f\alpha\\ &= \alpha'.\end{aligned}$$

□

Proposition 4.1. *Let (A, B, α) be a crossed module and (φ, X, ω) a lifting of (A, B, α). If (φ', X', ω') is a lifting of (A, X, φ) then $(\varphi', X', \omega\omega')$ is also a lifting of (A, B, α).*

Proof. The proof is immediate. □

Let (φ, X, ω) and (φ', X', ω') be two liftings of (A, B, α). A morphism f from (φ, X, ω) to (φ', X, ω') is a group homomorphism $f\colon X \to X'$ such that $f\varphi = \varphi'$ and $\omega' f = \omega$. Hence lifting crossed modules of (A, B, α) and morphisms between them form a category which we denote by $\mathsf{LXMod}/(A, B, \alpha)$. By Proposition 4.1 it follows that if φ is a lifting of (A, B, α) over $\omega\colon X \to B$ then $\mathsf{LXMod}/(A, X, \varphi)$ is a full subcategory of $\mathsf{LXMod}/(A, B, \alpha)$.

Let (A, B, α) be a transitive crossed module. Then a lifting (φ, X, ω) of (A, B, α) is called an *n-lifting* when $|\operatorname{Ker}\omega| = n$.

Corollary 4.1. *If (φ, X, ω) is a 1-lifting of (A, B, α) then ω is an isomorphism. Hence $(A, X, \varphi) \cong (A, B, \alpha)$.*

Proof. If (φ, X, ω) is a 1-lifting of (A, B, α), then ω becomes surjective and $|\operatorname{Ker}\omega| = 1$, i.e., ω is injective. Hence ω is an isomorphism. □

Theorem 4.3. *Let $(f, g)\colon (\widetilde{A}, \widetilde{B}, \widetilde{\alpha}) \to (A, B, \alpha)$ be a morphism of crossed modules where $(\widetilde{A}, \widetilde{B}, \widetilde{\alpha})$ is transitive and let (φ, X, ω) be a lifting of (A, B, α). Then there is a unique morphism of crossed modules $(f, \widetilde{g})\colon (\widetilde{A}, \widetilde{B}, \widetilde{\alpha}) \to (A, X, \varphi)$ such that $\omega\widetilde{g} = g$ if and only if $f(\operatorname{Ker}\widetilde{\alpha}) \subseteq \operatorname{Ker}\varphi$.*

Proof. Assume that $f(\operatorname{Ker}\widetilde{\alpha}) \subseteq \operatorname{Ker}\varphi$. For the existence; let $\widetilde{b} \in \widetilde{B}$. Since $(\widetilde{A}, \widetilde{B}, \widetilde{\alpha})$ is transitive, $\widetilde{\alpha}$ is surjective and there exists an $\widetilde{a} \in \widetilde{A}$ such that $\widetilde{\alpha}(\widetilde{a}) = \widetilde{b}$. Hence $\widetilde{g}(\widetilde{b}) = \varphi f(\widetilde{a})$.

It is easy to see that $\widetilde{g}$ is well defined since $f(\operatorname{Ker}\widetilde{\alpha}) \subseteq \operatorname{Ker}\varphi$. Also

$$\begin{aligned}\omega\widetilde{g}(\widetilde{b}) &= \omega\varphi f(\widetilde{a}) \\ &= \alpha f(\widetilde{a}) \\ &= g\widetilde{\alpha}(\widetilde{a}) \\ &= g(\widetilde{b}).\end{aligned}$$

So $\omega\widetilde{g} = g$. By the definition of $\widetilde{g}$ it implies that $(f, \widetilde{g})$ is a crossed module morphism. For any other morphism $g'\colon \widetilde{B} \to X$ such that $\omega g' = g$, the pair (f, g') is a crossed module morphism and coincides with $\widetilde{g}$.

Conversely suppose that $(f, \widetilde{g})\colon (\widetilde{A}, \widetilde{B}, \widetilde{\alpha}) \to (A, X, \varphi)$ is a crossed module morphism with $\omega\widetilde{g} = g$. If $\widetilde{a} \in \operatorname{Ker}\widetilde{\alpha}$, then $f(\widetilde{a}) \in f(\operatorname{Ker}\widetilde{\alpha})$ and $\widetilde{\alpha}(\widetilde{a}) = 0$. Since $(f, \widetilde{g})$ is a crossed module morphism $\widetilde{g}\widetilde{\alpha} = \varphi f$ and hence $\varphi f(\widetilde{a}) = 0$. So $f(\widetilde{a}) \in \operatorname{Ker}\varphi$ and hence $f(\operatorname{Ker}\widetilde{\alpha}) \subseteq \operatorname{Ker}\varphi$. □

Corollary 4.2. *Let (A, B, α) be a crossed module. Assume that (φ, X, ω) and $(\widetilde{\varphi}, \widetilde{X}, \widetilde{\omega})$ are two liftings of (A, B, α) such that $(A, \widetilde{X}, \widetilde{\varphi})$ is transitive. Then $\widetilde{\varphi}$ is a lifting of φ if and only if* $\operatorname{Ker}\widetilde{\varphi} \subseteq \operatorname{Ker}\varphi$.

Corollary 4.3. *Let (A, B, α) be a crossed module. Assume that (φ, X, ω) and $(\widetilde{\varphi}, \widetilde{X}, \widetilde{\omega})$ are two liftings of (A, B, α) such that (A, X, φ) and $(A, \widetilde{X}, \widetilde{\varphi})$ are both transitive. Then $(\varphi, X, \omega) \cong (\widetilde{\varphi}, \widetilde{X}, \widetilde{\omega})$ if and only if* $\operatorname{Ker}\varphi = \operatorname{Ker}\widetilde{\varphi}$.

Theorem 4.4. *Let (A, B, α) be a crossed module, X a group and let $\omega\colon X \to B$ be an injective group morphism. Then any group morphism $\varphi\colon A \to X$ such that $\omega\varphi = \alpha$ becomes a lifting of α over ω.*

Proof. According to Theorem 4.2 we only need to show that $\overline{\varphi}\colon A \ltimes X \to X$ defined by $\overline{\varphi}(a, x) = \varphi(a) + x$ is a group morphism, i.e. $\varphi(x \cdot a) = x + \varphi(a) - x$ for all $x \in X$, $a \in A$.

$$\begin{aligned} \omega(\varphi(x \cdot a)) &= \omega(\varphi(\omega(x) \cdot a)) \\ &= \alpha(\omega(x) \cdot a) && \text{(since } \omega\varphi = \alpha\text{)} \\ &= \omega(x) + \alpha(a) - \omega(x) && \text{(by CM1)} \end{aligned}$$

and

$$\begin{aligned} \omega(x + \varphi(a) - x) &= \omega(x) + \omega(\varphi(a)) - \omega(x) \\ &= \omega(x) + \alpha(a) - \omega(x) && \text{(since } \omega\varphi = \alpha\text{.)} \end{aligned}$$

Therefore since ω is injective and $\omega(\varphi(x \cdot a)) = \omega(x + \varphi(a) - x)$ then $\varphi(x \cdot a) = x + \varphi(a) - x$ for all $x \in X$, $a \in A$. This completes the proof. □

Corollary 4.4. *Every crossed module (A, B, α) lifts to the crossed module $(A, A/N, p)$ over $\omega\colon A/N \to B$, $a + N \mapsto \alpha(a)$ where $N = \operatorname{Ker}\alpha$.*



This lifting is called *natural lifting*. By Corollary 4.4 and first isomorphism theorem for groups we can say that for every crossed module (A, B, α), $(A, \operatorname{Im}\alpha, \alpha)$ is a transitive crossed module with the same action.

Now we will give a criterion for the existence of the lifting crossed module.

Theorem 4.5. *Let (A, B, α) be a crossed module and C a subgroup of $\operatorname{Ker}\alpha$. Then there exists a lifting (φ, X, ω) of α such that $\operatorname{Ker}\varphi = C$. Moreover in this case $\operatorname{Ker}\omega = \operatorname{Ker}\alpha/C$.*

Proof. Since $\operatorname{Ker}\alpha$ is central, C is a normal subgroup of A. By Corollary 4.4 $(\varphi, A/C, \omega)$ is a lifting of α where $\varphi(a) = a + C$ and $\omega(a + C) = \alpha(a)$. Obviously $\operatorname{Ker}\varphi = C$. Since φ is surjective $(A, A/C, \varphi)$ is transitive. Further

$$\begin{aligned}\operatorname{Ker}\omega &= \{a + C \mid \omega(a + C) = 0\}\\ &= \{a + C \mid \alpha(a) = 0\}\\ &= \{a + C \mid a \in \operatorname{Ker}\alpha\}\\ &= \operatorname{Ker}\alpha/C\end{aligned}$$

and this completes the proof. □

Definition 4.2. Let (A, B, α) be a crossed module. A lifting (φ, X, ω) of (A, B, α) is called *transitive* if both crossed modules (A, B, α) and (A, X, φ) are transitive.

Corollary 4.5. *Let (φ, X, ω) be a transitive lifting of (A, B, α). Then (φ, X, ω) is an n-lifting if and only if $|\operatorname{Ker}\alpha/\operatorname{Ker}\varphi| = n$.*

Corollary 4.6. *Let X be a topological group and C a subgroup of the fundamental group $\pi_1(X, 0)$ of X at the identity. Then there exists a lifting $(p, \mathsf{St}_{\pi X}0/C, \omega)$ of $(\mathsf{St}_{\pi X}0, X, d_1)$ such that $\operatorname{Ker}p = C$. Moreover if the underlying space of X is path-connected, then $\mathsf{St}_{\pi X}0/\pi_1(X, 0) \cong X$ as groups.*

Corollary 4.7. *If X is a topological group whose underlying space is totally disconnected, then $\mathsf{St}_{\pi X}0 \cong \pi_1(X, 0)$ as groups.*

Proposition 4.2. *If (φ, X, ω) is a 1-transitive lifting of a crossed module (A, B, α), then (φ, X, ω) is a lifting of any lifting of (A, B, α).*

Proof. Since (A, X, φ) is 1-transitive, $\operatorname{Ker}\varphi$ is trivial and hence for any lifting $(\widetilde{\varphi}, \widetilde{X}, \widetilde{\omega})$ the kernel $\operatorname{Ker}\varphi$ is contained in $\operatorname{Ker}\widetilde{\varphi}$. Further by Corollary 4.2 (φ, X, ω) is a lifting of $(\widetilde{\varphi}, \widetilde{X}, \widetilde{\omega})$. □

Hence one can state the following definition.

Definition 4.3. A lifting of a crossed module (A, B, α) is called *universal* if it lifts to every lifting of (A, B, α).

By Proposition 4.2 a 1-transitive lifting is universal.
As a consequence of Theorem 4.5 we can give the following corollary.

Corollary 4.8. *Every crossed module has a universal lifting.*

Proof. Let (A, B, α) be a crossed module. If we choose C as trivial in Theorem 4.5, then we have that $(1_A, A, \alpha)$ is a lifting of (A, B, α). Moreover $(1_A, A, \alpha)$ is a lifting of every lifting of (A, B, α). □

5 Equivalences of the categories

In this section we prove that for a certain group-groupoid G, there is a categorical equivalence between the group-groupoid actions of G on groups and lifting crossed modules of the crossed module corresponding to G. We also prove that the liftings of a crossed module are categorically equivalent to the covering morphisms of the same crossed module.

Theorem 5.1. *Let G be a group-groupoid and (A, B, α) the crossed module corresponding to G. Then the category* $\mathsf{GpGpdAct}(\mathsf{G})$ *of group-groupoid actions of G on groups and the category* $\mathsf{LXMod}/(A, B, \alpha)$ *of lifting crossed modules of (A, B, α) are equivalent.*

Proof. Define a functor $\theta\colon \mathsf{GpGpdAct}(\mathsf{G}) \to \mathsf{LXMod}/(A, B, \alpha)$ assigning each object (X, ω) of $\mathsf{GpGpdAct}(\mathsf{G})$ to a lifting (φ, X, ω) of (A, B, α) where

$$\varphi\colon A \to X, a \mapsto a \bullet 0_X$$

Conversely define a functor $\psi\colon \mathsf{LXMod}/(A, B, \alpha) \to \mathsf{GpGpdAct}(\mathsf{G})$ assigning each lifting (φ, X, ω) of (A, B, α) to a group-groupoid action (X, ω) of G on the group X via an action map defined by

$$G_{d_0} \times_\omega X \to X, (g, x) \mapsto g \bullet x = \varphi(g - 1_{d_0(g)}) + x.$$

Now prove that $\theta \circ \psi$ and $\psi \circ \theta$ are respectively naturally isomorphic to the identity functors based on the categories $\mathsf{LXMod}/(A, B, \alpha)$ and $\mathsf{GpGpdAct}(\mathsf{G})$. If (φ, X, ω) is a lifting of (A, B, α), then $(\theta \circ \psi)(\varphi, X, \omega) = (\varphi', X, \omega)$ where φ' is defined by

$$\varphi'(a) = a \bullet 0_X = \varphi(a - 1_{d_0(a)}) + 0_X = \varphi(a)$$

for all $a \in A$ and hence $\theta \circ \psi = 1$.

On the other hand if (X, ω) is an object of $\mathsf{GpGpdAct}(\mathsf{G})$ with an action of G on X given by

$$G_{d_0} \times_\omega X \to X, (g, x) \mapsto g \bullet x$$

then the new action obtained from the functor ψ is defined by

$$\begin{aligned} g \bullet' x &= (g - 1_{d_0(g)}) \bullet (0_X + x) \\ &= (g - 1_{d_0(g)}) \bullet (0_X + (1_{\omega(x)}) \bullet x) \\ &= (g - 1_{d_0(g)} + 1_{\omega(x)}) \bullet (0_X + x) \qquad (1_{d_0(g)} = 1_{\omega(x)}) \\ &= g \bullet x. \end{aligned}$$

and therefore $\psi \circ \theta = 1$ which completes the proof. □

Recall that as a result of [6, Proposition 4.2] a morphism $(f_1, f_2)\colon (\widetilde{A}, \widetilde{B}, \widetilde{\alpha}) \to (A, B, \alpha)$ of crossed modules such that $f_1\colon \widetilde{A} \to A$ an isomorphism is called *covering morphism*. Hence we can give the following theorem.

Theorem 5.2. *For a crossed module (A, B, α), the category $\mathsf{LXMod}/(A, B, \alpha)$ of lifting crossed modules and the category $\mathsf{CovXMod}/(\mathsf{A}, \mathsf{B}, \alpha)$ of covering crossed modules of (A, B, α) are equivalent.*

Proof. Clearly if (φ, X, ω) is a lifting of (A, B, α), then $(1_A, \omega)\colon (A, X, \varphi) \to (A, B, \alpha)$ is a crossed module morphism. Hence (A, X, φ) is a covering crossed module of (A, B, α). Moreover if $(\widetilde{\varphi}, \widetilde{X}, \widetilde{\omega})$ is another lifting of (A, B, α) and f is a morphism in $\mathsf{LXMod}/(A, B, \alpha)$ from (φ, X, ω) to $(\widetilde{\varphi}, \widetilde{X}, \widetilde{\omega})$, then $(1_A, f)$ is a morphism in $\mathsf{CovXMod}/(\mathsf{A}, \mathsf{B}, \alpha)$ from (A, X, φ) to $(A, \widetilde{X}, \widetilde{\varphi})$. This construction is clearly functorial.

Conversely if $(f_2, f_1)\colon (\widetilde{A}, \widetilde{B}, \widetilde{\alpha}) \to (A, B, \alpha)$ is a covering morphism of crossed modules, then $f_2\colon \widetilde{A} \to A$ is an isomorphism and $\varphi = \widetilde{\alpha} f_2^{-1}$ is the lifting of α over f_1. It is easy to see that $f_1\varphi = \alpha$ and $(A, \widetilde{B}, \varphi)$ is a crossed module. Further if

$$(g_2, G)\colon (A', B', \alpha') \to (A, B, \alpha)$$

is another covering of (A, B, α) and $(h_2, h_1)\colon (\widetilde{A}, \widetilde{B}, \widetilde{\alpha}) \to (A', B', \alpha')$ is a morphism in $\mathsf{CovXMod}/(\mathsf{A}, \mathsf{B}, \alpha)$ then $h_1\colon (\varphi, \widetilde{B}, f_1) \to (\varphi', B', G)$ is a morphism in $\mathsf{LXMod}/(A, B, \alpha)$. This construction is also functorial and the other details of the equivalence is straightforward. □

By Example 4.1 and Theorem 5.2 we can state the following corollary.

Corollary 5.1. *Every crossed module (A, B, α) is a covering of the automorphism crossed module $(A, \mathrm{Aut}(A), \iota)$ constructed by A.*

The category $\mathsf{CovXMod}/(\mathsf{A}, \mathsf{B}, \alpha)$ of crossed module coverings of a certain crossed module (A, B, α) is equivalent to the category $\mathsf{GpGpdCov}/\mathsf{G}$ of group-groupoid coverings of group-groupoid corresponding to (A, B, α) (see also [1, 15] for more general case of crossed modules over groups with operations for an algebraic category C). Hence by Theorem 5.2 we can obtain the following corollary.

Corollary 5.2. *Let G be a group-groupoid and (A, B, α) the corresponding crossed module. Then the category* $\mathsf{GpGpdCov/G}$ *of group-groupoid coverings of G and the category* $\mathsf{LXMod}/(A, B, \alpha)$ *of lifting crossed modules of (A, B, α) are equivalent.*

By Corollary 4.8 and Corollary 5.2 we can obtain the following result.

Corollary 5.3. *Every group-groupoid has a universal covering group-groupoid.*

By Corollary 4.4 and Corollary 5.2 we can obtain the following result.

Corollary 5.4. *Let G be a group-groupoid. Then G acts on $X = \operatorname{Ker} d_0/G(0)$ over $\omega\colon \operatorname{Ker} d_0/G(0) \to \operatorname{Ob}(G), \omega(a + G(0)) = d_1(a)$ as a group-groupoid action with*

$$\varphi\colon G_{d_0} \times_\omega X \to X, \varphi(b, a + G(0)) = (b \circ a) + G(0)$$

In [6, Proposition 2.3] it was proved that for a topological group X whose underlying space has a universal cover, the category $\mathsf{TGrCov/X}$ of topological coverings of X and the category $\mathsf{GpGpdCov}/\pi\mathsf{X}$ of coverings of group-groupoid πX are equivalent; and in [1, Theorem 4.1] a similar result was given for more general topological groups with operations. Hence we can state the following corollary.

Corollary 5.5. *Let X be a topological group whose underlying space has a universal cover. Then the category* $\mathsf{TGrCov/X}$ *of covers of X in the category of topological groups and the category*

$$\mathsf{LXMod}/(\mathsf{St}_{\pi X}0, X, d_1)$$

of liftings of $(\mathsf{St}_{\pi X}0, X, d_1)$ are equivalent.

6 Conclusion

Using the results of the paper [16] it could be possible to develop normal and quotient notions of lifting crossed modules. Moreover using the equivalence of the categories in [19, Section 3] as stated in the introduction, parallel results of this paper could be obtained for crossed modules of group with operations and internal groupoids working in a certain algebraic category C.

References

[1] H.F. Akız, N. Alemdar, O. Mucuk, T. Şahan, *Coverings of internal groupoids and crossed modules in the category of groups with operations*, Georgian Math. Journal, **20 (2)** (2013), 223–238.

[2] R. Brown, *Topology and Groupoids*, BookSurge LLC, North Carolina, 2006.

[3] J.C. Baez, A.D. Lauda, *Higher-dimensional algebra V: 2-groups* Theory Appl. Categ. **12** (2004), 423–491.

[4] R. Brown, P.J. Higgins, R. Sivera, *Nonabelian Algebraic Topology: filtered spaces, crossed complexes, cubical homotopy groupoids*, European Mathematical Society Tracts in Mathematics 15, 2011.

[5] R. Brown, G. Danesh-Naruie, J.P.L. Hardy, *Topological Groupoids: II. Covering Morphisms and G-Spaces*, Math. Nachr. **74** (1976), 143–156.

[6] R. Brown, O.Mucuk, *Covering groups of non-connected topological groups revisited*, Math. Proc. Camb. Phil. Soc. **115** (1994), 97–110.

[7] R. Brown, C.B. Spencer, *G-groupoids, crossed modules and the fundamental groupoid of a topological group*, Proc. Konn. Ned. Akad. v. Wet. **79** (1976), 296–302.

[8] T. Datuashvili, *Cohomologically trivial internal categories in categories of groups with operations*, Appl. Categ. Structures **3** (1995), 221–237.

[9] T. Datuashvili, *Cohomology of internal categories in categories of groups with operations, Categorical Topology and its Relation to Analysis, algebra and combinatorics*, Ed. J. Adamek and S. Mac Lane (Prague, 1988), World Sci. Publishing, Teaneck, NJ, 1989.

[10] T. Datuashvili, *Kan extensions of internal functors: Nonconnected case*, J.Pure Appl.Algebra **167** (2002), 195–202.

[11] T. Datuashvili, *Whitehead homotopy equivalence and internal category equivalence of crossed modules in categories of groups with operations*, Proc. A. Razmadze Math.Inst. **113** (1995), 3–30.

[12] P.J. Higgins, *Categories and groupoids*, Van Nostrand, New York, 1971.

[13] J.-L. Loday, *Spaces with finitely many non-trivial homotopy groups*, J. Pure Appl. Algebra **24** (1982), 179–202.

[14] O. Mucuk, B. Kılıçarslan, T. Şahan, N. Alemdar, *Group-groupoids and monodromy groupoids* Topology Appl. **158 (15)** (2011), 2034–2042.

[15] O. Mucuk, T. Şahan, *Coverings and crossed modules of topological groups with operations*, Turk. J. Math. **38 (5)** (2014), 833–845.

[16] O. Mucuk, T. Şahan, N. Alemdar, *Normality and Quotients in Crossed Modules and Group-groupoids*, Appl. Categor. Struct. **23** (2015), 415–428.

[17] O. Mucuk, H.F. Akız, *Monodromy groupoids of an internal groupoid in topological groups with operations*, Filomat **29 (10)** (2015), 2355–2366.

[18] G. Orzech, *Obstruction theory in algebraic categories I, II.* J. Pure Appl. Algebra **2** (1972), 287–314, 315–340.

[19] T. Porter, *Extensions, crossed modules and internal categories in categories of groups with operations*, Proc. Edinb. Math. Soc. **30** (1987), 373–381.

[20] J.H.C. Whitehead, *Note on a previous paper entitled "On adding relations to homotopy group"*. Ann. of Math. **47** (1946), 806–810.

[21] J.H.C. Whitehead, *On operators in relative homotopy groups*, Ann. of Math. **49** (1948), 610–640.

[22] J.H.C. Whitehead, *Combinatorial homotopy II*, Bull. Amer. Math. Soc. **55** (1949), 453–496.